



Necrotizing Colitis in an Adult Following SARS-CoV-2 Infection: A Case Report and Literature Review

**Mahmoud DABBAGH ^{a,b}, ELAZZAOUI Imad ^{a,b*},
Amine MAAZOUZ ^{a,b}, Hind HABLADJE ^{a,b}, Walid CHAIR ^{a,b},
Rania ELMOUHIB ^{a,b}, Houda BOUIDIDA ^{a,b},
Mehdi KARAMI ^{a,b}, Mohammed LAMGHARI ^{a,b},
Mohamed NAJIH ^{a,b}, Mountassir MOUJAHID ^{a,b},
Hakim ELKAOUI ^{a,b}, Ahmed BOUNAIM ^{a,b}
and Sidi Mohammed BOUCHENTOUF ^{a,b}**

^a Department of Visceral Surgery I, Military Hospital Mohamed V, Rabat, Morocco.

^b Faculty of Medicine and Pharmacy, University Mohamed V, Rabat, Morocco.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/127153>

Case Report

Received: 15/09/2024

Accepted: 19/11/2024

Published: 28/11/2024

ABSTRACT

Necrotizing colitis is a rare but life-threatening condition, often overlooked in young patients due to its uncommon occurrence. It is typically associated with premature neonates but can also affect adults, particularly those with vascular compromise or thrombotic conditions.

*Corresponding author: E-mail: imad.elazzaoui@gmail.com.

Cite as: DABBAGH, Mahmoud, ELAZZAOUI Imad, Amine MAAZOUZ, Hind HABLADJE, Walid CHAIR, Rania ELMOUHIB, Houda BOUIDIDA, Mehdi KARAMI, Mohammed LAMGHARI, Mohamed NAJIH, Mountassir MOUJAHID, Hakim ELKAOUI, Ahmed BOUNAIM, and Sidi Mohammed BOUCHENTOUF. 2024. "Necrotizing Colitis in an Adult Following SARS-CoV-2 Infection: A Case Report and Literature Review". *Asian Journal of Case Reports in Surgery* 7 (2):537-40. <https://journalajcrs.com/index.php/AJCRS/article/view/581>.

We describe the case of a 44-year-old male who was admitted for treatment of necrosis in the sigmoid colon caused by thrombosis in the inferior mesenteric artery, possibly triggered by a previous SARS-CoV-2 infection.

This case highlights the importance of considering SARS-CoV-2 as a potential trigger for thromboembolic events and the urgent need for multidisciplinary management in cases of necrotizing colitis.

Keywords: Necrotizing colitis; emergency surgery; thrombosis of the inferior mesenteric artery.

1. INTRODUCTION

Necrotizing colitis is an acute condition predominantly impacting premature neonates, characterized by considerable morbidity and mortality rates [1]. Its occurrence in adults is notably lower, with decreased mortality rates [2]. Prognosis relies mainly on urgent, multidisciplinary treatment.

2. CASE PRESENTATION

A 44-year-old young man with a history of SARS-Cov2 infection, admitted to the emergency department with a 4-day history of occlusion, clinical examination revealed a patient in relatively good general condition, with a heart rate of 110 bpm and blood pressure of 100/60 mmHg afebrile, the abdomen was distended with tenderness localized to the flank and left iliac fossa.

Biological screening showed hemoglobin level at 10g/dL, WBC 16400 elements/mm³, C-reactive protein 120 mg/L.

Abdominal CT scan with IV contrast showed necrosis of sigmoid colon due to thrombosis of inferior mesenteric artery with peritoneal fluid effusion.




Fig. 1. CT-Scan Showing Sigmoid Necrosis

Emergency surgery was performed, surgical exploration revealed localized necrosis of the sigmoid colon, which was subsequently resected with the confection of a double stoma.

Medical treatment was then instituted with curative-dose LMWH.

The follow-up was marked by a clinical and biological recovery. Patient was then referred to internal medicine department for etiological investigation.

3. DISCUSSION

"Necrotizing colitis (NC) predominantly impacts premature neonates of low birth weight, with an incidence ranging from 0.3 to 2.4 per 1000 live births, but notably higher at approximately 10% among infants weighing less than 1500 grams" [3,4]. "While NC occurrence is substantially lower in adults and carries a decreased mortality rate, it still presents significant morbidity" [5].

"Several causes can lead to NC in adults. Factors such as infectious agents, inflammatory mediators, and circulatory disturbances are all implicated in its etiology and pathogenesis. A wide range of infectious bacterial causes can lead to NC, including *Klebsiella*, *E. coli*, *Clostridia*, and *Staphylococcus epidermidis*, as well as viruses like Coronaviruses, Rotaviruses" [2].

The clinical manifestations of NC are variable and depend mainly on the severity and time course of the disease. Main symptoms are abdominal pain, vomiting and nausea, abdominal tenderness, abdominal contracture or tenderness, bowel obstruction.

"Patients rarely have fever and won't have hyperleukocytosis. In cases of severe intestinal ischemia causing transmural infarction and necrosis, abdominal wall contracture may be noted on abdominal examination and is often accompanied by metabolic acidosis and shock" [6].




Fig. 2. Image showing the necrotic sigmoid colon

“Ultrasonography and radiography can be used as diagnostic tools, but a CT scan has been declared to be the first choice for the diagnosis of NC [7]. The images that can be seen are intestinal wall thickening, per colon fat infiltration, with or without peritoneal fluid. Computed tomography may suggest a vascular etiology; in the case of complete obstruction, the intestinal wall will be thin and non-enhanced, with dilatation of the lumen, possibly with thrombosis in the mesenteric vessels” [8].

“Overall, a CT scan can be used to determine the severity and exact location of the injury. Also, MR angiography can be used to evaluate any suspected arteriovenous IMA malformation” [7].

“The innumerable causes of colonic ischemia can be categorized into occlusive and nonocclusive pathologies. Thrombophilia and vasculitis are the most important among the occlusive causes of ischemic colitis in the young” [9].

SARS-CoV-2 has been linked to the development of coagulopathy and thromboembolic complications in severe COVID-19 patients. The presentations included pulmonary embolism, deep venous thrombosis, and rarely mesenteric ischemia. Our patient had history of SARS-CoV-2 infection.

“Treatment is multidisciplinary and involves surgery and LMWH-based medical therapy. Surgical intervention is necessary if there is intestinal necrosis or frank perforation or when there is clinical deterioration over 12–24 hours

despite intensive medical support, surgical options are either resection of the necrotic segment with anastomosis, or resection and provision of a stoma” [2].

4. CONCLUSION

NC very rarely affects adults, its clinical presentation is widely variable, and its diagnosis is based on CT scans.

Management is urgent and relies mainly on surgery and administration of LMWH.

CONSENT

As per international standard or university standard, patient written consent has been collected and preserved by the authors.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the authors.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Fick KA, Wolken AP. NECROTIC JEJUNITIS. Lancet. 1949;253(6552):519-521.
DOI: 10.1016/S0140-6736(49)90847-8
2. Katara AN, Chandiramani VA, Soman R, Bhaduri A, Desai DC. Necrotizing enterocolitis in adults: A study of four cases. Indian J Surg. 2004;66.
Available: <https://tspace.library.utoronto.ca/handle/1807/2845>
Accessed May 4, 2024.
3. Chung DH, Ethridge RT, Kim S, et al. Molecular mechanisms contributing to necrotizing enterocolitis. Ann Surg. 2001; 233(6):835-842.
DOI: 10.1097/00000658-200106000-00014
4. Huang FS, Warner BW. Necrotizing enterocolitis: A closer look. Gastroenterology. 2001;121(4):1016-1017.
DOI: 10.1016/S0016-5085(01)70081-9
5. Mackay WD, Dawson JL. Necrotizing colitis. Proc R Soc Med. 1967;60(5):502.
DOI: 10.1177/003591576706000527
6. Washington C, Carmichael J. Management of Ischemic Colitis. Clin Colon Rectal Surg. 2012;25(4):228.
DOI: 10.1055/S-0032-1329534
7. Seeliger B, Agnus V, Mascagni P, et al. Simultaneous computer-assisted assessment of mucosal and serosal perfusion in a model of segmental colonic ischemia. Surg Endosc. 2020;34(11):4818-4827.
DOI:10.1007/S00464-019-07258-Z/METRICS
8. Case Study: Non - Embolistic Necrotizing Colitis - An Examination of Clinical Presentation and Management, IJSR, Call for Papers, Online Journal.
Available: <https://www.ijsr.net/getabstract.php?paperid=SR23727135155>
Accessed May 12, 2024.
9. Doulberis M, Panagopoulos P, Scherz S, Dellaporta E, Kouklakis G. Update on ischemic colitis: from etiopathology to treatment including patients of intensive care unit. Scand J Gastroenterol. 2016; 51(8):893-902.
DOI: 10.3109/00365521.2016.1162325

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/127153>