



Hemipteran Insect Pests Associated with Different Agricultural Crops from Chhatrapati Sambhajnagar (M.S), India

Muley E. D ^{a*}, Chavan A. S ^a and Naphade S. R ^a

^a Department of Zoology, Dr. Babasaheb Ambedkar Marathwada University, Chhatrapati Sambhajnagar- 431 004, Maharashtra, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.56557/ajorib/2025/v7i1105>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://prh.globalpresshub.com/review-history/1785>

Original Research Article

Received: 24/10/2024

Accepted: 27/12/2024

Published: 15/01/2025

ABSTRACT

A field investigation was conducted in Chhatrapati Sambhajnagar, Maharashtra, from May 2023 to April 2024 to look into the seasonal occurrence of hemipteran pests in different agricultural sectors. In this study, 10 hemipteran insect species from 9 genera, 6 families, and 8 subfamilies were collected from the study area. The Aphididae family was the most abundant. This study aimed to give a brief overview of hemipteran insects and the damages they cause to agricultural crops in the Chhatrapati Sambhajnagar district.

Keywords: Seasonal incidence; hemipteran insects; damages; agricultural crops.

*Corresponding author: Email: muleyeshwar@gmail.com;

1. INTRODUCTION

Insects are among the most prevalent creatures to visit crop fields, acting as pests, pollinators, and other ecologically important organisms to a particular plant (Mitra et al., 2018). The global diversity of tropical insect fauna significantly exceeds predictions, as insects are key components of animal diversity in terms of the number of species in most habitats and ecosystems (Stork, 1988). Insect pests and illnesses, which thrive in intensive agriculture, are significant barriers to higher yield. Resistance, more outbreaks, and new pests have resulted in increased financial losses. Pest infestation reduces the quality and quantity of agricultural crops. According to Singh et al., (2003), insect diseases and pests can account for up to 38% of agricultural losses in India, followed by weeds at 10% and birds at 1-2%. Timely identification of agricultural pests is essential for a successful integrated pest management (IPM) program.

Hemipteran insects, usually known as 'true bugs', are economically significant because the majority of them are pests of a variety of agricultural crops. Hemiptera is classified as a monophyletic group because they have unique piercing and sucking mouth parts called rostrums, which are made up of concentric stylets that interlock to produce the feeding and salivary canal (Backus, 1988). Not all Hemipteran insects are a nuisance. Furthermore, numerous hemipterans are essential natural predators of several insect pests.

The world's known Hemipteran species number 103,590 and are classified into 152 families and four suborders. According to ZSI (ZSI, 2012) and Chandra (Chandra, 2011), the Indian Hemipteran fauna consists of 6479 species distributed across 92 families. The impact of hemipteran insects on agricultural crops in the Chh. Sambhajinagar (M.S) region of India has become an issue of worry for both agricultural practitioners and researchers. Many pestiferous and predatory insect species of the order Hemiptera have been incompletely reported for agricultural crops in this district. Therefore, the current study aims to identify the hemipteran species associated with diverse agricultural fields in the Chhatrapati Sambhajinagar district of Maharashtra, India. The purpose of this study is to provide insight into the incidence of hemipteran insects on agricultural crops and how they may affect crop productivity. By

gathering this information, researchers and farmers may work together to develop sustainable and ecologically friendly pest management solutions adapted to local conditions, assuring the long-term health and productivity of agricultural crops in the area.

2. MATERIALS AND METHODS

This study was conducted in Chh. Sambhajinagar district of Maharashtra, located at 19.8776° N 75.3423° E. Based on the convenience of season and cropping patterns, different sites from Chh. Sambhajinagar were selected for the study of the infection of hemipteran bugs associated with different agricultural crops. The adult bugs were collected from six different sites (Satara tanda, Deolai, Gandheli, Sindon, Bhindon, Dr. B. A. M. University Campus) during May 2023 to April 2024. Numerous agricultural sites from the study area were randomly visited for the collection of different methods such as the sweep net method, hand picking, and baiting were used for the collection of bugs. Specimens collected from various agricultural crop fields were killed in killing bottles (containing chloroform / ethyl acetate), pinned, preserved dry, and labelled to indicate host plant, locality, and date. During the present study, the standard methodology suggested by Borror and White (Borror & White, 1970), (Richards & Davies, 1997), (Triplehorn & Johnson, 2005), was adopted for the collection, preservation, and identification of hemipteran insects.

3. RESULTS AND DISCUSSION

During the present investigation, Hemipteran species related with different agricultural crops and other plant species such as soyabean, pigeon pea, cotton, wheat, maize, mung bean, parthenium and mulberry etc., were observed and collected. A total of 10 species belonging to 7 families, 7 subfamilies and 09 different genera were collected from the study area. Among these families, Pentatomidae was the most diversified while Aphididae was most abundant (in case of frequency).

According to Graph 1, members of family Aphididae showed highest abundance (in case of frequency) while members of family Coreidae showed lowest frequency. (Aphididae> Pentatomidae>Scutelleridae>Reduviidae>Pyrrhocoridae>Plataspidae>Coreidae).




Photo Plate 2. Hemipteran Insects Collected from Chhatrapati Sambhajinagar District


Table 1. Taxonomic Composition of hemipteran insects from Chhatrapati Sambhajnagar

Order	Family	Subfamily	Scientific Name	Common Name
Hemiptera	Reduviidae	Ectrichodiinae	<i>Scadra annulipes</i> (Reuter, 1881)	Assassin bugs
	Plataspidae	Plataspinae	<i>Megacopta cribraria</i> (Fabricius, 1798)	Bean plataspid
	Pentatomidae	Pentatominae	<i>Halyomorpha halys</i> (Stal, 1855)	Shield bugs
			<i>Nezara viridula</i> (Linnaeus, 1758)	Stink bugs
			<i>Dictyotus caenosus</i> (Westwood, 1837)	Brown shield bug
	Coreidae	Coreinae	<i>Anoplocnemis phasiana</i> (Fabricius, 1781)	Leaf footed bugs
	Pyrrhocoridae	Pyrrhocorinae	<i>Dysdercus cingulatus</i> (Fabricius, 1775)	Cotton stainer
	Scutelleridae	Scutellerinae	<i>Chrysocoris stollii</i> (Wolff, 1801)	Jewel bug
	Aphididae	Aphidinae	<i>Aphis gossypii</i> (Glover, 1877)	Cotton aphid
			<i>Aphis craccivora</i> (Koch, 1854)	Cowpea aphid

Table 2. Damage symptoms of Hemipteran insects on different host plants

Sr. No	Name of the Hemipteran Species	Host Plants	Damage Symptoms
1.	<i>Scadra annulipes</i>	Cotton, Pigeon pea	-Sucks fluids from leaves & stems -Discolouration, yellowing -Some assassin bugs can be beneficial as natural enemies of pests.
2.	<i>Megacopta cribraria</i>	Pigeon pea, soyabean, cotton, wheat, citrus, mulberry	-Feeds on underside of the leaves and stems -Abnormal plant development -Formation of necrotic areas
3.	<i>Halyomorpha halys</i>	Soyabean, citrus, maize	-Sucks plant sap -Adults feed on fruits -Lesions, necrotic areas on leaves
4.	<i>Nezara viridula</i>	Soyabean, mung bean, pigeon pea, cow pea	-Feeds on plant sap -Brownish spots on fruits -Retarded fruit growth
5.	<i>Dictyotus caenosus</i>	Soyabean, mung bean, cotton	-Feed on developing seed -Reduced seed quality
6.	<i>Anoplocnemis phasiana</i>	Cow pea, mung bean	-Adults suck the juice of unripe seeds -tender part gets dried

Sr. No	Name of the Hemipteran Species	Host Plants	Damage Symptoms
7.	<i>Dysdercus cingulatus</i>	Cotton, okra	-Sucks fluid from host plants -Affects the flower, seed capsule/ boll, Staining of cotton bolls
8.	<i>Chrysocoris stollii</i>	Pigeon pea, mung bean, soyabean	-Feed on plant sap -Adults feed on leaves, seeds and inflorescence
9.	<i>Aphis gossypii</i>	Cotton, citrus, pigeon pea, soyabean, okra	-Adults feed on underside of leaves -Curling of leaves, Foliage may die permanently
10.	<i>Aphis craccivora</i>	Cow pea, pigeon pea, soyabean	-Sucks plant sap -Pods may become malformed -Drying of affected pods



Graph 1. Frequency of Hemipteran Insects Collected from Chh. Sambhajinagar

Similar studies regarding various aspects of hemipteran insects including their diversity and damaging effects were carried out by numerous researchers. Kandibane, (2008) conducted studies on diversity and abundance of phytophage hemipteran insects in an irrigated rice ecosystem of Tamil Nadu and recorded 24 species of hemipteran insects. Kailash Chandra and Kushwaha (2013) studied the hemipteran insect pest diversity associated with *Lablab purpureus* L. from Jabalpur, Madhya Pradesh and recorded 7 species of hemipteran insects. Burman and Gupta [(2016) observed the assemblage of coleoptera and hemiptera community in Chakrashila Wildlife Sanctuary in Assam and collected 3 species of hemipteran insects. Aseefa and Malindzisa (2018) carried out a molecular identification of mealybugs on cultivated, ornamental and wild host plants in Swaziland and revealed the presence of six mealybug species belonging to four genera. Harshita et al., (2022) carried out a study on the seasonal incidence of major insect pests of vegetable pigeon pea (*Cajanus cajan* L.) and concluded that *N. viridula* was one of the most severe pests of pigeon pea crop.

Sahoo et al., (2021) conducted a study on the diversity and abundance of Hemipteran families at Agri-biodiversity park of Prof. Jayashankar Telangana State Agricultural University, Hyderabad. They recorded a total of 12,575 individuals under 22 families of Hemiptera. Seni (2021) carried out a study on the hemipteran insect fauna associated with pigeon pea and their relative abundance at Chiplima, Odisha and documented a total of 18 hemipteran insect taxa belonging to 11 insect families. Paunikar et al., (2022) studied the diversity and distribution of hemipteran fauna in the different forest areas of North-West Himalaya and collected 27 species of 23 genera under 7 families, belonging to order Hemiptera. All these works indicate that severe infestations of insect pests can affect the yield of agricultural, horticultural crops and also affect the normal growth of various other plants species.

4. CONCLUSION

The present investigation made an effort to provide an overview of Hemipteran insects in the Chhatrapati Sambhajanagar district. The results of this investigation will help us understand the diversity of hemipteran insects and their infestations on diverse agricultural crops in the Chhatrapati Sambhajanagar district. A long-term study is required to investigate species

distribution across all seasons as well as how they interact with host plants as the environment changes. In addition to providing baseline data for future researchers and a large possibility for additional research, it provides useful information regarding the diversity of hemipteran insects in the area mentioned.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology.

Details of the AI usage are given below:

1. Authors hereby declare that generative AI technologies such as Quill Bot software has been used during the writing or editing of manuscripts for grammar checking and summarizing.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Amerika Singh, A. S., Saroj Singh, S. S., & Rao, S. N. (2003). Integrated pest management in India. In *Integrated pest management in the global arena* (pp. 209-221). Wallingford, UK: CABI Publishing.
- Assefa, Y., & Malindzisa, N. (2018). Molecular identification of mealybugs (Hemiptera: Pseudococcidae) on cultivated, ornamental, and wild host plants in Swaziland. *Asian Research Journal of Agriculture*, 9, 1-10.
- Backus, E. A. (1988). Sensory systems and behaviours which mediate hemipteran plant-feeding: A taxonomic overview. *Journal of Insect Physiology*, 34(3), 151-165. <https://doi.org/xxxxxxx>
- Barman, B., & Gupta, S. (2016). Assemblage of coleoptera and hemiptera community in a stream of Chakrashila Wildlife Sanctuary in Assam. *Tropical Ecology*, 57(2), 243-253.
- Borror, D. J., & White, R. E. (1970). *A field guide to insects: America north of Mexico*. Houghton Mifflin Harcourt.

- Chandra, K. (2011). Insect fauna of states and union territories in India. *ENVIS Bulletin: Wildlife & Protected Areas*, 14(1), 189-218.
- Chandra, K., & Kushwaha, S. (2013). Record of hemipteran insect pest diversity on *Lablab purpureus* L: An economically important plant from Jabalpur, Madhya Pradesh. *Research Journal of Agricultural Sciences*, 4(1), 66-69.
- Harshita, A. P., Gopali, J. B., Ramanagouda, S. H., Basavaraja, N., Rudresh, D. L., & Mulla, S. R. (2022). Seasonal incidence of major insect pests infesting vegetable pigeon pea. *Pharma Innovation Journal*, 23-26.
- Kandibane, M. (2008). Diversity and abundance of phytophage hemiptera in an irrigated rice ecosystem of Tamil Nadu, India. *Asian Journal of Bioscience*, 3(1), 102-106.
- Mitra, B., Shah, S. K., & Mishra, P. (2018). Insect fauna associated with the tea ecosystem of North Bengal, India. *Records of the Zoological Survey of India*, 178-193.
- Paunekar, S. D., Kushwaha, S., Saha, P. C., & Jahan, S. (2022). Diversity and distribution of some hemiptera fauna in the different forest areas of North-West Himalaya. *Journal of Natural Resource & Development*, 17(2), 75-82.
- Richards, O. W., & Davies, R. G. (1997). *Imms' general textbook of entomology* (10th ed.). New Delhi: B I Publication Pvt. Ltd.
- Sahoo, K. C., Sunitha, V., Rao, V. V., & Chary, D. S. (2021). Diversity of hemipteran families at Agri-biodiversity park, Hyderabad, India. *Entomon*, 46(2), 143-148.
- Seni, A. (2021). Diversity of hemipteran insect fauna and their relative abundance in pigeon pea, *Cajanus cajan* (L.) Millsp. *Journal of Entomology and Zoology Studies*, 9(1), 1072-1079.
- Stork, N. E. (1988). Insect diversity: Facts, fiction and speculation. *Biological Journal of the Linnean Society*, 35(4), 321-337. <https://doi.org/xxxxxxx>
- Triplehorn, C. A., & Johnson, N. F. (2005). *Borror and DeLong's introduction to the study of insects*. Brooks/Cole.
- ZSI. (2012). COP XI Publications. Retrieved from <http://www.zsi.gov.in/Cop-11/cop-11.html>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://prh.globalpresshub.com/review-history/1785>