

Frequency of HCV and HBV Co-infections in HIV Positive Patient in City of Iran: A Cross-Sectional Study

**Fatemeh Saleh¹, Hossein Azizi², Farnaz Kheirandish^{3*}, Fariborz Rashnou⁴,
Seyed Dawood Mousavi Nasab⁵, Fatollah Movahedi⁶ and Maryam Azizi⁷**

¹Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran, Expert Laboratory Shohadaye Ashayer Hospital, Khorram Abad, Iran.

²Department of Microbiology, Borujerd Branch, Islamic Azad University, Borujerd, Iran, Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran.

³Department of Parasitology and Mycology, Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran.

⁴Department of Surgery, Lorestan University of Medical Sciences, Khorramabad, Iran.

⁵Department of Microbiology, Borujerd Branch, Islamic Azad University, Borujerd, Iran.

⁶Lorestan University of Medical Sciences, Khorramabad, Iran, Expert Laboratory Shohadaye Ashayer Hospital, Khorram Abad, Iran.

⁷Department of Microbiology, Yazd Branch, Islamic Azad University, Yazd, Iran.

Authors' contributions

This work was carried out in collaboration between all authors. Author FS did the study design and wrote the protocol. Authors FS and HA and FKH did the statistical analysis and literature searches while analyses of study was by authors FR, FM and MA. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJTDH/2015/13282

Editor(s):

(1) Amr M. Mohamed, Department of Animal Medicine, Assiut University, Egypt.

(2) William Ebomoyi, Department of Health Studies, College of Health Sciences, Chicago State University, USA.

Reviewers:

(1) Anonymous, Sweden.

(2) Luiz Euribel Prestes Carneiro, Infectology Department, Regional Hospital and University of Oeste Paulista, Presidente Prudente, Brazil.

(3) Maurice Y. Mongkuo, Government & History, Fayetteville State University, USA.

(4) Anonymous, India.

(5) Anonymous, Iran.

(6) Anonymous, Congo.

Complete Peer review History: <http://www.sciencedomain.org/review-history.php?iid=851&id=19&aid=7432>

Original Research Article

Received 11th August 2014
Accepted 8th December 2014
Published 19th December 2014

ABSTRACT

Aims: One of the current crises of the society is prevalence rate of viral co-infection of hepatitis B and C among HIV patients. HIV, HBV, and HCV are major public health concerns. Because of shared routes of transmission, HIV-HCV coinfection and HIV-HBV coinfection are common. HIV-positive persons are at risk of being infected with HBV and HCV. Frequency of co-infection with HBV and HCV in HIV-patients is depended on the geographic regions and the type of exposure. The aim of this study was to determine the prevalence of HIV, HBV and HCV co-infection in one of the medical educational centers of Lorestan province, Iran.

Study Design: This Cross-sectional One Shot Case Study Design to determine the prevalence of concurrent viral infection in the city of Khorramabad, Iran.

Place and Duration of Study: Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran and Shohadaye Ashayer hospital, Khorramabad, Iran, in 2013

Methodology: This cross-sectional study was designed in 2013 in order to study the prevalence of co-infection among 500 outpatients and inpatients referring to Shohadaye Ashayer Hospital, Khorramabad, Lorestan province. The existence of specific antigen and antibody against the virus was determined by Enzyme-Linked Immunosorbent Assay. (Demographic information was extracted from the patients' medical files and frequency of three viruses was calculated according to demographics information.

Results: 103 out of 500 studied serum samples were infected with HBV, HCV, and HIV viruses. Maximum infection was in the age range of more than 40 years old, which was equal to 23 patients (24.7%). 58 out of 103 samples were inpatients, 18 (9.4%) of whom were co-infected. 3 out of 103 studied samples (5.9%) who were co-infected with HIV/HBV/HCV were male. On the other hand, 4 (3.9%) male patients were infected with HIV/HBV and 4 (7.8%) male patients were co-infected with HBV/HCV. Also, 23 (22.3%) patients were co-infected with HIV/HCV; one (1.9%) was female and 22 (43.1%) were male.

Conclusion: Patients who have both hepatitis B and C infections when are associated with HIV infection the disease becomes more severe. By obtain frequency, relationship was detected between hospital care of patients and prevalence of con-infection. Due to frequent injections in the hospital and taking blood and its products, inpatients were more susceptible to infection than outpatients. So, the probability of their infection with viruses was higher than outpatients. This issue is a disturbing factor in medicinal centers which should receive more attention.

Keywords: Co-infection; hepatitis B; hepatitis C; ELISA; HIV.

1. INTRODUCTION

HIV, hepatitis C virus (HCV), and hepatitis B virus (HBV) are three important factors of blood-borne diseases [1]. Hepatitis B is one of the most prevalent infectious diseases in the world. About 400 million worldwide are carriers of hepatitis B virus. According to the recent report of World Health Organization, after tuberculosis and malaria, hepatitis B is the most prevalent contagious and infectious disease with the annual emergence of 50 million people [2]. These patients are also exposed to the risk of liver cirrhosis and hepatocellular carcinoma (HCC). Prevalence of hepatitis B carriers in Iran is 3% of the society or about two million people [3]. HCV infection is another blood-borne infection which has infected approximately 170 million people in the world [3]. Hepatitis C is among the important factors for chronic liver

diseases and, due to its long-term treatment process, can lead to cirrhosis and liver cancers [4]. Infection with HCV and HBV are two important factors of mortality and disability, which impose financial expenses to the society and are among the important problems of world health [4]. AIDS is a spectrum of abnormalities caused by disorder in the function of cellular and humoral immunity which is resulted from HIV infection [4]. According to the recent statistics, about 33 million HIV-infected patients are currently living in the world and more than 20 million of them have died so far. 5800 people per day are infected with this virus, 90% of whom are in developing countries [4]. In addition to blood transfusion, this virus can be transferred via other ways such as injection of intravenous drugs by infected instruments, taking coagulating factors, hemodialysis, homosexual and heterosexual contacts, and from mother-to-child-

transmission (MTCT) [5,6]. According to Screening of all donated blood for hepatitis B, which already takes place and increasing use of hepatitis B virus vaccine, fortunately, the prevalence record of this infection has been remarkably decreased. But, due to the lack of a vaccine for preventing the infections caused by hepatitis C and HIV viruses, infection with these diseases has still remained a challenge. Fortunately, in Iran the prevalence of HIV among blood donors is not very high and it shows blood safety [7,8]. Patients hospitalized in different medical centers, especially dialysis, thalassemia, and hemophilia patients, are more liable than outpatients to infection with viral infections and consequently hospital infections, which is due to frequent receiving of blood products and coagulated concentrates [9,10]. AIDS affects all the body systems due to causing extreme immune disorders. These disorders have been comprehensively investigated in the world. Mortality risk of liver diseases has a reverse relationship with the number of CD₄s [11]. Increased rate of hepatocellular carcinoma hepatotoxicity due to taking antiretroviral drugs has been also observed among the patients co-infected with HCV, HBV, and/or HIV [11,12]. Considering the accompanying problems of HIV patients, the probability of infection with hepatitis B and C is very high among them [13]. Transfer risk of infectious diseases through blood can be measured by analyzing and studying information of blood donors, screening methods, and prevalence rate of serologic markers of infective diseases [13]. Obviously, the important point about these infections is determining the prevalence rate of carriers and patients in the region in order to plan and adopt effective strategies in terms of prevention and control by the related organizations and develop an appropriate ground for researchers and thus assist medical and health systems using the statistic data obtained from relative frequency distribution of these infections and their analysis [14]. So, the goal of the study was to examine the co-frequency of these transferable infections among inpatients and outpatients referring to Shohadaye Ashayer Hospital, Khorramabad, Iran, in 2013.

2. METHODOLOGY

This cross-sectional study was carried out from March to February 2013. 500 serum samples were collected from outpatients and inpatients referring to Shohadaye Ashayer Hospital, Khorramabad, Lorestan province. Samples were

studied to investigate the presence of HIV, HBV, and HCV viruses antigens using ELISA kit (past kit, Tehran, Iran- pishtazteb, Tehran, Iran-Aesku/ISA. Diagnostics, Germany) and Biotech kit (Ireland) for antibody. Sampling was based on the request of the physician and patient satisfaction.

Studying the files of the participants led to obtaining some information such as age, gender, and inpatient or outpatient conditions. ELISA reader device (Stat Fax, model 2100, USA) was used to read the results of ELISA test.

3. RESULTS

Out of 500 patients, 103 cases were infected with HCV, HBV, and HIV viruses, 52 were female and 51 were male, respectively. Age range was between 20 and 40 years old. 5% of patients were below 20 years and among whom only 1 case (1.1%) had HIV-HCV co-infection. Maximum infection was in the age range of more than 40, which was equal to 23 people (24.7%). Out of 103 samples, 58 specimens were related to inpatients, 18 people (19.4%) of whom had co-infection (Fig. 1). Out of 103 studied samples, 20 samples were infected with HIV, 10 people (19.2%) of whom were female and 10 (19.6%) were male. Seventeen patients were only infected with HBV, 12 people (23.1%) of whom were female and 5 (9.8%) were male. Thirty-two patients were HCV positive, 29 people (55.8%) of whom were female and 3 (5.9%) were male (Table 1).

Out of 103 studied samples, 3 specimens (2.9%) had HIV/HBV/HCV co-infection, all of whom were male. Also 4 (3.9%) male patients were infected with HBV/HIV and 4 (7.8%) male patients had HBV/HCV co-infection. Moreover, 23 (22.3%) patients had HIV/HCV co-infection, among whom 1 (1.9%) was female and 22 (43.1%) were male. According to the frequency of infection in Hospitalized patients, was found that, hospitalization of patients can be a factor for the generation of co-infection, specifically in patients with impaired immune system.

4. DISCUSSION

One of the most important and prevalent diseases accompanying HIV infection is hepatitis B and C and different studies have been done in this regard. When the patients with both types of hepatitis B and C are co-infected with HIV, they will suffer from more severe complications [15].

According to the obtained results, men had more 33 of men were infected, while just 1 female patient had this co-infection. In a study at Imam Khomeini Hospital, Tehran, Iran, in 2005– 2010, 85 (29.2%) out of 291 male patients had HIV/HBV/HCV co-infection and, in comparison with the present study, the prevalence of co-infection among the studied male patients was higher during a year [11]. In a research in the west of Iran, maximum rate of infection was in the average age range of 31-50 years old, which was in accordant with the current study, and can be due to the point that hepatitis vaccination of children which was started from 1993 in Iran somehow decreased such infection among the youth in the west of Iran [12]. In a study in 2011, maximum rate of co-infection with these three types of viruses was reported in the age range of less than 40 at Tehran AIDS Research Center and between 21-46 years old at Imam Khomeini Hospital, which was somehow contrary to the co-infection prevalence in the present investigation [11,13]. Also, in a study in Tehran by Rahimi A and et al. [14], maximum co-infection was in the age range of less than 20 years old. A study in Brazil showed that maximum viral co-infection (HIV/HCV) was among young men (71%) [15]. Since Iran is an Islamic country and people follow Islamic religious principles, less prevalence of viral infections among the youth has been reported in this country. In western societies, a higher degree of infection has been reported due to free homosexual and heterosexual contacts [15]. Among the inpatients of this study, about 19.4% were co-infected with HIV/HBC/HCV, which may indicate that while patients are hospitalized due to infection with one of these viruses, they simultaneously suffer from other underlying viral diseases. Moreover, due to frequent injections and intake of blood

HIV/HBV/HCV co-infection than women so that and its products, inpatients are more exposed to infections than outpatients; therefore, they are in a worst condition and have higher risk of infection with other viral diseases compared with outpatients [16,17]. Also in this study all tests were repeated twice to ensure accuracy. The molecular tests such as PCR is not performed in the hospital, after positive tests, PCR to patients in other centers such as blood transfer is recommended. Considering the above-mentioned points and comparison of the frequency of co-infection in this study with other investigations, and the number of patients and study period, it was found that the prevalence of these infections was considerable in Lorestan province. This issue can be due to different reasons such as high degree of unemployment which was about 12.2% in this province [18]. Unemployment can cause several problems such as sociological problems, increased psychological abnormalities, and addiction which may result in increased infections. Psychological abnormalities, addiction, and lack of knowledge about infection manner and other cases are among the factors that highly influence increased prevalence of these viral diseases [18,19]. Finally, it is recommended to take some measures for freely and confidentially testing the patients with high risk. This issue can be influential in the decrease of viral infections, especially co-infection, and improvement of the society's health [20]. This study aimed to examine HBV and HCV coinfection serologically and determine the shared and significant factors in the coinfection of HIV-positive patients was conducted. Provide different information from different geographic locations can make decisions in the field of infection control and effective global public health.




Fig. 1. Studying co-infection prevalence in patients with HIV/HBV/HCV co-infection in the age range of less than 20 to more than 40 in Shohadaye Ashayer Hospital, Khorramabad, Lorestan province, Iran

Table 1. Prevalence of co-infection HIV-HBV-HCV in male and female patients admitted to Shohadaye Ashayer hospital, Khorramabad, Iran, in 2013

Genus	HIV	HCV	HBV	HIV-HCV	HIV-HBV	HCV-HBV	HIV-HBV-HCV	Total
Male	10(19.6%)	3(5.9%)	5(9.8%)	22(43.1%)	4(3.9%)	4(7.8%)	3(5.9%)	51
Female	10(19.2%)	29(55.8%)	12(23.1%)	1(1.9%)	0	0	0	52
Total	20(19.4%)	32(31.1%)	17(16.5%)	23(22.3%)	4(3.9%)	4(3.9%)	3(2.9%)	103

5. CONCLUSION

Out of 103 studied samples, 20 samples were infected with HIV, 10 people (19.2%) of whom were female and 10 (19.6%) were male. Seventeen patients were only infected with HBV, 12 people (23.1%) of whom were female and 5 (9.8%) were male. Thirty-two patients were HCV positive, 29 people (55.8%) of whom were female and 3 (5.9%) were male. This study aimed to examine HBV and HCV coinfection serologically and determine the shared and significant factors in the coinfection of HIV-positive patients was conducted.

CONSENT

All authors declare that 'written informed consent was obtained from the patient (or other approved parties) for publication of this case report and accompanying images.

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki. The study was approved by the Medical Research Ethics Committee of Lorestan University of Medical Sciences with a letter to the Issue 129/93/6012.

ACKNOWLEDGMENTS

The authors would like to express their gratitude to the respectable head and colleagues at Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, for their sincere cooperation. We also appreciate all the people who helped us in this project.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Adewole OO, Anteyi E, Ajuwon Z, Wada I, Elegba F, Ahmed P, et al. Hepatitis B and C virus co-infection in Nigerian patients with HIV infection. *J Infect Dev Ctries.* 2009;3(5):369-75.
2. Pourazar A, Akbari N, Hariri M, Yavari F, Akbari Sh. Evaluation of demographic Profiles and Prevalence of major Viral markers in first time vs repeat blood donors in Esfahan. *SJIBTO.* 2006;2(7):323-329.
3. Jaiswal SP, Chitnis DS, Naik G, Artwani KK, Pandit CS. Prevalence of Anti-HCV in central India. *Indian J Med Res.* 1996;104:177-81.
4. 2013. Available: www.unaids.org/esources/campaigns/globalreport
5. Garfein RS, Vlahov D, Galai N, Doherty MC, Nelson KE. Viral infections in short-term injection drug users: The prevalence of the hepatitis C, hepatitis B, human immunodeficiency, and human T-lymphotropic viruses. *Am J Public Health.* 1996;86.
6. Hollinger FB, Purcell RH, Gerin JL. *Viral Hepatitis.* Philadelphia: Lippincott, Williams & Wilkins; 2002.
7. Okada S, Taketa K, Ishikawa T, Koji T, Swe T, Win N, et al. High prevalence of hepatitis C in patients with thalassemia and patients with liver diseases in Myanmar (Burma). *J Acta Med Okayama* 2000;54(3):137-8.
8. Ardalan FA, Osquei MRF, Toosi MN, Irvanloo G. Synergic effect of chronic hepatitis C infection and beta thalassemia major with marked hepatic iron overload on liver fibrosis: a retrospective cross-sectional study. *BMC Gastroenterol.* 2004;4:17.
9. Mannucci PM. Hemophilia and related bleeding disorders: A story of dismay and success. *Hematology. Am. Soc. Hemato. Educ. Program.* 2002;1-9.

10. Koziel MJ, Peters MG. Viral hepatitis in HIV infection. *N Engl J Med*. 2007;356(14):1445-1454.
11. Banafsheh Moradm, Badie, Seyed Ahmad Seyed Alinaghi, Reza Chaman, et al. Frequency and correlates of co-infection Hepatitis C and Hepatitis B with HIV. *Knowledge & Health*. 2011;6(3):40-43.
12. Mohsen Mohammadi, Gholamreza Talei, et al. Survey of both hepatitis B virus (HBsAg) and hepatitis C virus(HCV-Ab) coinfection among HIV positive patients. *Virology Journal*. 2009;6:202
13. Amitis Ramezani, Mohammad Banifaz, et al. Occult Hepatitis B virus infection: A major concern in HIV-infected patients Occult HBV in HIV. *Hepat Mon*. 1. 2011;11(1):7-10.
14. Afarin Rahimi-Movaghar, Emran M. Razaghi, Elaheh, Sahimi-Izadian, Masoumeh Amin-Esmaeili. HIV, Hepatitis C virus, and Hepatitis B virus co-infections among injecting drug users in Tehran, Iran. *International Journal of Infectious Diseases*. 2010;14:28-33.
15. Valdete Meurer Kuehlkamp, Fabiana Schuelter-Trevisol. Prevalence of human immunodeficiency virus/Hepatitis C virus co-infection in Brazil and associated factors: The Brazilian Journal of Infectious Diseases. 2013;17(4):455-463.
16. David K. Henderson. Managing occupational risks for Hepatitis C transmission in the health care setting. *Clin Microbiol Rev*. 2003;16(3):546-568.
17. Hadadi A, Haji Abdolbaghi M, Karbakhsh M, Rasoulinejad M, Mohraz M. The effectiveness of prophylaxis following occupational exposure to hepatitis B and HIV in medical professions. *Journal of Tropical Infectious Diseases*. 2007;12(38):33-38.
18. Available:<http://khabaronline.ir/detail/286609/economy/market>
19. Hadadi A, Afhami SH, Kharbakhsh M, Hajabdoulbaghi M, Rasoulinejad M, Emadi H, et al. Epidemiological determinants of occupational exposure to HIV, HBV, HCV, in health care workers. *Tehran University Medical Journal*. 2007;65(9):59-66.
20. Maryam Yarmohammadi. Investigating the serologic status and epidemiological aspects of health care workers' exposure to HBV and HCV viruses. *Knowledge & Health*. 2011;5(4):37-42.

© 2015 Saleh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history.php?iid=851&id=19&aid=7432>