



Clinicopathological Study of Pityriasis Alba in Libyan Patients

Jumana Eswaysi ^a, Aisha Trabelsi ^b, Ibrahim Almukahal ^b
and Gamal Duweb ^{b*}

^a Dermatology Department, Alwahda Teaching Hospital, Derna, Libya.

^b Dermatology Department, Faculty of Medicine, Benghazi university, Benghazi, Libya.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajrdes/2025/v8i1118>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129753>

Original Research Article

Received: 08/11/2024

Accepted: 14/01/2025

Published: 18/01/2025

ABSTRACT

Abstract: Pityriasis alba is a common, benign skin disorder occurring predominantly in children and adolescents. Most patients have a history of atopy, and pityriasis alba may be a minor manifestation of atopic dermatitis. It is characterized by ill-defined macules and patches or thin plaques, round or oval, often with mild scaling, and sometimes with mild pruritus. The lesions initially may be mildly erythematous, and over time they become hypopigmented. They are most commonly located on the face (especially the cheeks), arms, and upper trunk, and they are more noticeable in people with darker skin types. Sun exposure accentuates the lesions. Pityriasis alba resolves spontaneously, with a gradual return of normal skin pigmentation.

Aim of the Study: To assess the etiological and physical characteristics in Libyan pityriasis alba patients

Participants and Methods: One hundred patient attending to dermatology department outpatient clinic, Alwahda hospital-Derna-Libya are included in this descriptive study. Diagnosis of PA is done

*Corresponding author: E-mail: drduweb@gmail.com;

by dermatological clinical examination. Detailed history of age, gender ; history of atopy, sun exposure, frequency of bathing , with physical examination of site, number of lesion is recorded.

Results: The study included 100 Libyan pityriasis alba patient, 69(69%) male, 31 (31%) female, their age ranged from 2-19 years. 89% of patient's lesions were on face only , 8% on face and extremities and 3% on face and neck. The number of pityriasis alba lesions ranged from 2 to 12 lesions, scales were evident in only 73% of patients and 27% had no scaly lesions. The duration of the disease ranged from 1 week to 2 years. 66% were Fitzpatrick skin type IV and 34% skin type III. Xerosis presented in 70%. History of atopic dermatitis was +ve in 11% and -ve in the other 89%. History of atopy +ve in 55%, -ve in 45% of patients Family History of atopy + ve in 90% and -ve in 10% . In 37% of Pityriasis alba patients, the lesions were recurrent and 63% had no history of recurrence. Only 4% of our patients were using sunscreens, Sun exposure_hrs per day ranged between 1 to 6 hrs. The times of bathing per week ranged from 1 to 7 times per week.

Conclusion: Our data suggested that pityriasis alba is more prone in children of skin type IV with strong family history of atopy, and strong association with skin xerosis, Sun exposure and neglected sunscreen usage. To the best of our knowledge, this is the first publication about pityriasis alba from our country.

Keywords: Pityriasis alba; derna-libya; atopy; xerosis; fitzpatrick skin types.

1. INTRODUCTION

Pityriasis alba, a relatively common skin disorder in children and young adults, is characterized by the presence of ill-defined, scaly, faintly erythematous patches. These lesions eventually subside, leaving hypopigmented areas that then slowly return to normal pigmentation (see the images below)" (Miazek et al., 2015; Jadotte and Janniger, 2011). "Lesions may progress through the following three clinical stages: Papular (scaling) erythematous Papular (scaling) hypochromic Smooth hypochromic. No known cause of pityriasis alba has been reported. The condition is not contagious, and no infectious agent has been identified. Leading theories as to the origin of the lesions in pityriasis alba have involved atopy and post inflammatory changes". (Rao et al., 2023) "A large number of patients with pityriasis alba have a history of atopic disease, and atopic patients are prone to developing the condition" (Rao et al., 2023). "Other theories of origin include hypopigmentation secondary to pityriacitrin, a substance produced by Malassezia yeasts that acts as a natural sunscreen. In one study, atrophic sebaceous glands were noted in almost half the cases of pityriasis alba. In another study, iron deficiency anemia was reported in 16.5% of patients" (Vinod et al., 2002), however, "this may be a coincidental finding, and the clinical relevance of anemia is not yet known. Excessively dry skin, which is frequently exacerbated by cold, dry environments, also appears to be a common factor in pityriasis alba.

Reported contributory factors related to the development of pityriasis alba are excessive and

unprotected sun exposure, poor hygienic habits, and environmental influences such as temperature, humidity, and altitude" (Blessmann Weber et al., 2002). "In addition, one study found patients with pityriasis alba to be more likely to have low levels of serum copper, which could contribute to tyrosinase dysfunction and thus a decrease in the production of melanin" (Bechelli et al., 1981). "Although the exact incidence has not been described, as many as 5% of children may have pityriasis alba" (Plensdorf et al., 2017). "The disorder is not seasonal, but the dry, slightly scaly appearance tends to worsen during cold months, when the air is relatively dry inside the home. In addition, sun exposure may make the lesions more obvious during spring and summer. The condition is more common in patients with a history of atopy. In a study of 9955 school children aged 6-16 years who lived in a tropical region, the prevalence of pityriasis alba was 9.9%" (Carneiro et al., 2014). In a study from South India, the overall frequency of hypopigmentary disorders among children seen in a dermatology outpatient department was 3.28 per 1000 (Weber et al., 2002). In the 113 children treated for hypopigmentary disorders over a 23-month study period, pityriasis alba was the most common diagnosis (24.7%). "Another study, in Nepal, showed that the prevalence of pityriasis alba within a wide range of dermatoses was 5.2%" (Bassaly et al., 1963). "Age-, sex-, and race-related demographics Pityriasis alba is most common in children aged 3-16 years, with 90% of cases occurring in children younger than 12 years. The disease occasionally is found in adults" (Elesawy et al., 2020). "Males and females are equally susceptible to the disease,

but a slight male predominance has been noted" (Jadotte and Janniger, 2011; Blessmann Weber et al., 2002). "Pityriasis alba occurs in people of all races. In one study, the incidence of PA was slightly high in light-skinned people, whereas other studies have demonstrated a higher incidence among more darker skin" (Blessmann Weber et al., 2002; Ofarrell, 1956). "The condition is frequently more apparent and cosmetically bothersome in patients with darker complexions" (Blessmann Weber et al., 2002; Ofarrell, 1956). "Some scholars believe that the occurrence of the disease may be related to personal hygiene habits such as skin dryness caused by regular bathing, etc" (Weber et al., 2002). "There are 2 clinical variants of classic PA (CPA): endemic PA, which occurs in children living in poor socioeconomic conditions, and atopy-related" (Ruiz-Maldonado, 2007). The improvement of endemic PA in a few months with mild topical hydrating creams, topical antimicrobials, and sunscreens, and its deterioration with the use of topical corticosteroids and calcineurin inhibitors, suggest distinctive etiologies (Ruiz-Maldonado, 2007), with clear prognostic and therapeutic implications.

"Classic PA also must be distinguished of Extensive Pityriasis Alba (EPA). Both show fewer melanocytes and melanosomes" (Martin et al., 1990) and "both manifest as hypopigmented macules with scaling. However, EPA is more common in adults, has a more generalized and symmetric distribution, tends to appear on the trunk, and usually is not associated with a history of atopy or evidence of erythema" (Lin and Janniger, 2005).

2. AIM OF THE STUDY

The study aim was to assess the etiological and physical characters of PA patients in Derna - Libya.

3. MATERIALS AND METHODS

A total of 100 Libyan patient with pityriasis alba attending the outpatients department of Alwahda teaching hospital, Derna - Libya, were included in this cross-sectional study. Each patient exposed to detailed history and complete dermatological examination according to the prepared proforma as following:

1. Detailed disease history to record age, sex, duration of lesions, history of recurrence, history

of atopic dermatitis, atopy and family history of atopy, times of bathing, number of sun exposure and sunscreen usage.

2. Dermatological examination of all patients was carried out in order to determine skin type, xerosis, number of lesions and the site involved.

3.1 Statistical Analysis

Data were analyzed using statistical package for social science (SPSS) version 22. Descriptive analysis, frequencies, mean, standard deviation (SD) and chi square used for analysis.

4. RESULTS

The study included 100 Libyan pityriasis alba patients. Sixty nine patients (69%) were males and 31patients (31%) were females (Table 1), The age ranged from 2-19 years with median of 8.2 years and mean $\pm$ SD: 7.50 $\pm$ 3.45 years (Fig. 1).

In 89% of patients lesions were on face only, 8% on face and extremities and 3% on face and neck (Fig. 2). The number of pityriasis alba lesions ranged from 2 to 12 lesions with mean $\pm$ SD of 4.53 $\pm$ 1.9, 31% of patients had 4 to 7 lesions 2 to 3 lesions in 32% and more than 7 lesions in 7% of patients (Fig. 3). Scales were evident in only 73% of patients and 27% had no scaly lesions. The duration ranged from 1 week to 2 years with mean $\pm$ sd 15.34 $\pm$ 14.22736 months (Fig. 4). According to physical characteristics of participant patients , 66% were Fitzpatrick skin type IV and 34% skin type III (Fig. 5). Xerosis presented in 70% and 30% were clear (Fig. 6). In the study patients, history of atopic dermatitis was positive in 11% and history of atopy was positive in 55%. Family history of atopy was recorded in 90%. In 37% of PA patients the lesions were recurrent. only 4% of our patients were using sunscreens. Sun exposure ranged between __1 to 6 hrs with mean of 2.79 $\pm$ SD 1.391 hrs per day , 56% with more than 2hrs per day and 44% 2hrs or less (Fig. 7).The times of bathing per week ranged from 1 to 7 times (Fig. 8).

5. DISCUSSION

"Pityriasis Alba is a common skin disorder affecting mainly children and young adults; however, its etiopathogenesis is poorly understood, and there are many discrepancies about its Epidemiology in the literature. So this study is designed to assess the possible risk

Table 1. Gender distribution across groups

Group	Gender		Total
	Male	Female	
Case group	69 69.0%	31 31.0%	100 100.0%
Control group	9 36.0%	16 64.0%	25 100.0%
Total	78 62.4%	47 37.6%	125 100.0%

Pearson Chi-Square 9.283, $P = 0.002$ (Significant)




Fig. 1. Age parameters comparison across the study groups




Fig. 2. Distribution of the study participants with pityriasis alba according to the anatomical sites of the lesions




Fig. 3. Distribution of the study participants with pityriasis alba according to the number of the lesions




Fig. 4. Distribution of the study participants with pityriasis alba according to the duration of the lesion




Fig. 5. Distribution of the study participants with pityriasis alba according to the Fitzpatrick skin type




Fig. 6. Distribution of the study participants with pityriasis alba according to the presence of xerosis




Fig. 7. Distribution of the study participants with pityriasis alba according to the duration of daily sun exposure




Fig. 8. Distribution of the study participants with pityriasis alba according to the habit of bathing per week

factors of pityriasis alba. The present study described the clinical characteristics of the diagnosed lesions of PA among patients whereas their number range: 2 – 12 lesions, mean $\pm$ SD: 4.53 ± 1.9 and they were commonly small in size with scales in 73% of patients. and more than one third were recurrent. Some literature described the disease as rounded, oval or irregular plaque with indistinct margins” (Plensdorf et al., 2017; Novembre et al., 2001; Miazek et al., 2015). “Other mentioned that PA lesions are asymptomatic hypo-pigmented patches on the face, neck, trunk and proximal extremities of children. The single lesion has sharply demarcated margins and is covered by a fine branny scale” (Lee et al., 2018). “Moreover, it was reported that most lesions persist for some months and some may still show hypopigmentation for a year or more after all scaling subsides. Recurrent crops of new lesions may develop at intervals. The average duration of the common facial form in childhood is a year or more. as in our study lesion persist more than a year in 38% of patients. Pityriasis alba is found almost entirely in preadolescent children. In most instances, the lesions clear at puberty, however persistence into adulthood has been reported” (Ofarrell, 1956; Miazek et al., 2015; Lee et al., 2018). This study showed that the age of patients ranged from 2-19 years with median of 8.2 years and mean 7.50 This is slightly different from many reports at the age of 3–16 years old (Jadotte and Janniger, 2011) and mean of 9.6 years old (Plensdorf et al., 2017; Rao et al., 2023; Elshafey et al., 2012). “Differences in age distribution of PA may be due to differences in culture, ethnic group and external environment. Both sexes are equally susceptible to the disease, but a slight male predominance has been noted” (Plensdorf et al., 2017; Vinod et al., 2002; Blessmann Weber et al., 2002). “A clinico epidermiological study was carried out in 200 patients attending the skin department of H Command hospital, Air Force, Bangalore by Vinod et al The study group comprised of 115 (57.5%) males and 85 (42.5%) females” (Rao et al., 2023). The male to female ratio was 1.35:1 whereas in our study group which comprised of 100 PA patients (69%) males and (31%) females with ratio of 2.2 : 1 . A study carried out by Martin RF et al., (1990) and other studies found that PA has equal incidence in both boys and girls. In other hand extensive pityriasis alba a clinical variant of PA has female predominance (Martin et al., 1990; Rao et al., 2023).

“Regarding sun exposure, in study done by Khafagy et al. (2020) they found that it increased the risk of PA by more than four times. Our study is compatible with this because 56% of our patients exposed to sun for more than two hours per day and 96% are not using sun screens This may be explained by the fact that sun exposure causes tanning of the surrounding skin, which makes affected areas more prominent. This finding may give the reason why the majority of lesions occur in the face as our study found that 89% of PA lesion are on face only .and Vinod et al face was the only site involved in 91% of patients” (Plensdorf et al., 2017; Rao et al., 2023; Vinod et al., 2002) , “Essentially, PA lesions are more observable in darker- skinned patients. In this study, we found that pa was more prevalent in skin photo type IV and III (66%skin phototype IV and 44 % skin phototype III). These results were consistent with the study conducted by Ysabel et al.and other authors found that skin phototype IV was most frequently affected (61.8%) and skin phototype III (31.3%)” (Chang and Vashi, 2016, Godse et al., 2021, Bechelli et al., 1981) and Khafagy found skin phototype III or IV increases the risk of PA by more than four times (Khafagy et al., 2020). Elshafey et al. found that PA was more prevalent in individuals with darker skin and higher phototype categories (74.8% of patients with types VI and V and Blessmann et al., (2002) who revealed that “ultraviolet light B (UVB) had a strong influence on the development of PA in dark-skinned individuals (OR = (21,23), In our study, there was significant relation between PA and patients personal hygiene assessed by asking them about the frequency of bathing per week which showed that more than 70% of patients take a bath more than three times weekly . other studies found an increased incidence of PA among those with more frequent bathing” (Weber et al., 2002). “In this study there was no significant relation between pityriasis alba and the frequency of bathing, This may be due to removal of normal epidermal defensins and other natural protective substances from the skin surface by frequent bathing making one more prone to have pityriasis alba. On the contrary, Elshafey et al found no relation between the incidence of pityriasis alba and frequency of bathing” (Elshafey et al., 2012). “Atopic diathesis has commonly been described as a pathogenic factor in PA and some authors have even stated that PA is a feature of a tendency towards depigmentation, found exclusively in atopic patients” (Vinod et al., 2002). “An unequivocal relationship between some signs of AD (xerosis,palmar hyperlinearity,

Daniel Morgan folds) and PA was found: Blessmann et al., found that the presence of atopy signs was more common in PA patients; furthermore, he found important linear correlation between PA and atopic dermatitis signs, specifically when considering the presence of two or three signs of atopy" (Blessmann Weber et al., 2002).

Although atopic symptoms are often absent at the onset of PA, they may appear later in life (Blessmann Weber et al., 2002; Abdel-Wahab and Ragaie, 2022) this agree with this study as xerosis was found in 70 % of patients.

6. CONCLUSION

Pityriasis alba constitute a major health issue, males were commonly affected, with mean age of 7.5 years. In majority of patients lesions were on face only and skin type IV. Personal and family history of atopy was high. The mean time of sun exposure was 2.79 hours/ day and there was no significant correlation with frequency of bathing.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

CONSENT AND ETHICAL APPROVAL

The study was approved by hospital ethical committee. Every participant was informed and they willing accepted participation in the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abdel-Wahab, H. M., & Ragaie, M. H. (2022). Pityriasis alba: Toward an effective treatment. *J Dermatolog Treat*, 33(4), 2285-2289.
- Bassaly, M., Miale, A., & Prasad, A. S. (1963). Studies on pityriasis alba: A common facial skin lesion in Egyptian children. *Arch Dermatol*, 88(3), 272.
- Bechelli, L. M., Haddad, N., Pimenta, W. P., et al. (1981). Epidemiological survey of skin diseases in schoolchildren living in the Purus Valley (Acre State, Amazonia, Brazil). *Dermatologica*, 163(1), 78-93.
- Blessmann Weber, M., Sponchiado de Avila, L. G., Albaneze, R., Magalhães de Oliveira, O. L., Sudhaus, B. D., & Cestari, T. F. (2002). Pityriasis alba: A study of pathogenic factors. *J Eur Acad Dermatol Venereol*, 16(5), 463-468.
- Blessmann Weber, M., Sponchiado de Avila, L. G., Albaneze, R., Magalhães de Oliveira, O. L., Sudhaus, B. D., & Ferreira Cestari, T. (2002). Pityriasis alba: A study of pathogenic factors. *J Eur Acad Dermatol Venereol*, 16(5), 463-468.
- Carneiro, F. R. O., Do Amaral, G. B., Mendes, M. D., et al. (2014). Tissue immunostaining for factor XIIIa in dermal dendrocytes of pityriasis alba skin lesions. *An Bras Dermatol*, 89(2), 245-248.
- Chang, Y. C., & Vashi, N. A. (2016). Pityriasis alba. In *Clinical Cases in Skin of Color* (pp. 169–176). Springer International Publishing.
- Elesawy, F. M., Akl, E. M., & Halim, W. A. A. (2020). Zinc has a role in pathogenesis of pityriasis alba. *Indian J Paediatr Dermatol*, 21(3), 178-183.
- Elshafey, W. S., Fiala, L. A., Mohamed, R. W., & Ismael, N. A. (2012). The distribution and determinants of pityriasis alba among elementary school students in Ismailia City. *Journal of American Science*, 8(12), 444-449.
- Godse, P. A., Ajmera, K., & Godse, G. (2021). Atopic dermatitis in skin of color. In *Atopic Dermat. Pruritus Interes. Cases* (pp. 76-78).
- Jadotte, Y. T., & Janniger, C. K. (2011). Pityriasis alba revisited: Perspectives on an enigmatic disorder of childhood. *Cutis*, 87(2), 66-72.
- Khafagy, G. M., Nada, H. R., Rashid, L. A., El-Samanoudy, S. I., & Abd El-Sattar, E. M. (2020). Role of trace elements in pityriasis alba. *J Trace Elem Med Biol*, 59, 126422.
- Lee, D., Kang, J., Kim, S., et al. (2018). Case of extensive pityriasis alba. *Ann Dermatol*, 3(20), 146-148.
- Lin, R. L., & Janniger, C. K. (2005). Pityriasis alba. *Cutis*, 76, 21-24.

- Martin, R. F., Lugo-Somolinos, A., & Sanchez, J. L. (1990). Clinicopathologic study on pityriasis alba. *Bol Asoc Med P R*, 82(10), 463-465.
- Miazeck, N., Michalek, I., Pawlowska-Kisiel, M., Olszewska, M., & Rudnicka, L. (2015). Pityriasis alba—Common disease, enigmatic entity: Up-to-date review of the literature. *Pediatr Dermatol*, 32(6), 786-911.
- Miazeck, N., Michalek, I., Pawlowska-Kisiel, M., Olszewska, M., & Rudnicka, L. (2015). Pityriasis alba—Common disease, enigmatic entity: Up-to-date review of the literature. *Pediatr Dermatol*, 32(6), 786-911.
- Novembre, E., Cianferoni, A., Lombardi, E., et al. (2001). Natural history of "intrinsic" atopic dermatitis. *Allergy*, 56, 452-453.
- O'Farrell, N. M. (1956). Pityriasis alba. *Arch Dermatol*, 73(4), 376-377.
- Plensdorf, S., Livieratos, M., & Dada, N. (2017). Pigmentation disorders: Diagnosis and management. *Am Fam Physician*, 96(12), 797-804.
- Plensdorf, S., Livieratos, M., & Dada, N. (2017). Pigmentation disorders: Diagnosis and management. *Am Fam Physician*, 96(12), 797-804.
- Rao, M., Young, K., Jackson-Cowan, L., et al. (2023). Post-inflammatory hypopigmentation: Review of the etiology, clinical manifestations, and treatment options. *J Clin Med*, 12(3).
- Rao, M., Young, K., Jackson-Cowan, L., Kourosh, A., & Theodosakis, N. (2023). Post-inflammatory hypopigmentation: Review of the etiology, clinical manifestations, and treatment options. *J Clin Med*, 12(3).
- Ruiz-Maldonado, R. (2007). Hypomelanotic conditions of the newborn and infant. *Dermatol Clin*, 25, 373-382, ix.
- Vinod, S., Singh, G., Dash, K., & Grover, S. (2002). Clinico epidemiological study of pityriasis alba. *Indian J Dermatol Venereol Leprol*, 68(6), 338-340.
- Vinod, S., Singh, G., Dash, K., & Grover, S. (2002). Clinico epidemiological study of pityriasis alba. *Indian J Dermatol Venereol Leprol*, 68, 338-340.
- Weber, M. B., de Avila, L. G. S., Albaneze, R., et al. (2002). Pityriasis alba: A study of pathogenic factors. *J Eur Acad Dermatol Venereol*, 16(5), 463-468.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/129753>